Page 1/11




FIG. 1


BEST AVAILABLE COPY

Page 2/11

Id: 5'CTGGGATCCATCGAGGGTAGGATGCTGAAAGAAAAGAAGTACTCCCCGTCCTCTC-3'
Ir: 5'GTACTCCCCGTCCTCTGAAAGACCGTCACGACTTCGATCCGGGCGACCCAG-3'
IId: 5'-GAAGTGTCTCCGCTGAAAGACCGTCACAGCTTCTAGAAGCTTGCGGC-3'
IIr: 5'-GCCGCAAGCTTCTAGAAGCTGTGACGGTCTTCAGCGGAGACACTTC-3'

FIG. 2

Page 3/11






FIG. 3 A and B

Page 4/11






FIG. 3C

Page 5/11

A



C



В




FIG. 4

Page 6/11




FIG. 5

Page 7/11

| Preparation | Composition of regular structure, % | |
|-----------------------------|-------------------------------------|-------------|
| | α-structure | β-structure |
| Albebetin _{teor} # | 30 | 36 |
| Albebetin | 29 | 40 |
| Albeferon | 27 | 35 |
| Albeferon-I | 30 | 29 |
| Albeferon-II | 23 | 38 |

^{*}Obtained from theoretical analysis of the pre-designed structure of albebetin.

FIG. 6

Page 8/11

| Preparation | Antiviral activity | | |
|---------------------------|----------------------------|----------------------------|--|
| | (minimal concentration, M) | | |
| | Culture L-41 | culture VERO | |
| Albeferon | 1.3±0.61×10 ⁻⁵ | 8.5±6.8×10 ⁻⁵ | |
| Albeferon-I | 2.9±1.36×10 ⁻⁸ | 1.5±0.73×10 ⁻⁸ | |
| Albeferon-II | 1.4±0.45×10 ⁻¹⁰ | 1.4±0.46×10 ⁻¹⁰ | |
| Interferon-α ₂ | 5.5±3.57×10 ⁻¹¹ | 1.1±0.36×10 ⁻¹¹ | |
| Interferon-γ | 1.1±0.51×10 ⁻¹⁰ | 1.5±0.92×10 ⁻⁹ | |

FIG. 7

Page 9/11




FIG. 8

Page 10/11

| Preparation | Cytotoxical dose (CTD ₅₀), M | |
|--------------|--|-----------------------|
| | culture L-41 | culture VERO |
| Albebetin | ≥8.7×10 ⁻⁵ | ≥8.5×10 ⁻⁵ |
| Albeferon | ≥8.8×10 ⁻⁵ | ≥8.8×10 ⁻⁵ |
| Albeferon-I | ≥8.9×10 ⁻⁵ | ≥9.0×10 ⁻⁵ |
| Albeferon-II | ≥1.4×10 ⁻⁴ | ≥1.2×10 ⁻⁴ |

FIG. 9

Page 11/11

| Preparation | Antiviral activity (minimal concentration, M) | |
|---------------------------|---|----------------------------|
| | | |
| | culture L-41 | culture VERO |
| Peptide LKEKKYSP | 1.2±0.44×10 ⁻⁵ | 7.2±5.9×10 ⁻⁵ |
| Peptide LKDRHDF | 3.1±1.27×10 ⁻⁸ | 1.8±0.81×10 ⁻⁸ |
| Peptide LKEKKYSP-Ser-Ser- | 1.2±0.43×10 ⁻¹⁰ | 1.3±0.51×10 ⁻¹⁰ |
| Interferon-α2 | 5.0±3.49×10 ⁻¹¹ | 1.3±0.33×10 ⁻¹¹ |

FIG. 10

This Page is Inserted by IFW Indexing and Scanning Operations and is not part of the Official Record

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

BLACK BORDERS

IMAGE CUT OFF AT TOP, BOTTOM OR SIDES

FADED TEXT OR DRAWING

BLURRED OR ILLEGIBLE TEXT OR DRAWING

SKEWED/SLANTED IMAGES

COLOR OR BLACK AND WHITE PHOTOGRAPHS

GRAY SCALE DOCUMENTS

LINES OR MARKS ON ORIGINAL DOCUMENT

REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY

IMAGES ARE BEST AVAILABLE COPY.

☐ OTHER:

As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.